



**UNIVERSITI PUTRA MALAYSIA**

**TIDAL INFLUENCE ON THE PHYSICO-CHEMICAL PARAMETERS,  
TEMPORAL AND SPATIAL DISTRIBUTION OF ZOOPLANKTON IN  
LANGAT RIVER-ESTUARINE AREA, MALAYSIA**

**R.P PRABATH KRISHANTHA JAYASINGHE.**

**FS 2005 44**

**TIDAL INFLUENCE ON THE PHYSICO-CHEMICAL PARAMETERS,  
TEMPORAL AND SPATIAL DISTRIBUTION OF ZOOPLANKTON IN  
LANGAT RIVER-ESTUARINE AREA, MALAYSIA**

**By**

**R.P. PRABATH KRISHANTHA JAYASINGHE**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in  
Fulfilment of the Requirements for the Degree of Master of Science**

**June 2005**



## **Dedication**

To my mother

For her endless efforts

To give us the best life.....

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in  
fulfilment of the requirement for the degree of Master of Science

**TIDAL INFLUENCE ON THE PHYSICO-CHEMICAL PARAMETERS,  
TEMPORAL AND SPATIAL DISTRIBUTION OF ZOOPLANKTON IN  
LANGAT RIVER-ESTUARINE AREA, MALAYSIA**

By

**R.P. PRABATH KRISHANTHA JAYASINGHE**

**June 2005**

**Chairperson: Professor Fatimah Md. Yusoff, PhD**

**Faculty: Science**

The objectives of present study were to study the water quality, zooplankton distribution and its community structure along a salinity gradient in Langat river estuarine system ( $2^{\circ} 46'N$ ,  $101^{\circ} 26' E$ ), during low and high tides. Five sampling stations from the coastal area to the upstream were chosen for this study. Station 1 was in the coastal area while Station 5 was at the most upstream, the rest of the Stations (Stations 2, 3 and 4) are located in between, passing along the river gradient. Zooplankton and water samples were collected and analyzed monthly for a period of 12 months (January to December 2003).

The water quality, nutrients and zooplankton distribution changed according to the stations and tides. The mean salinity ranged from  $32.94 \pm 1.20$  PSU in



coastal waters to  $27.78 \pm 1.21$  PSU in upstream station during high tide. During low tide, the mean salinity in the most upstream station was as low as  $1.00 \pm 0.90$  PSU. Higher Dissolved Oxygen values ranged (means of  $5.84 \pm 0.35$  to  $7.09 \pm 0.13$   $\text{mgL}^{-1}$ ) during high tide and it decreased (means of  $3.91 \pm 0.71$  to  $5.22 \pm 0.71$   $\text{mgL}^{-1}$ ) during low tide. On the other hand, the nutrients such as total ammonia nitrogen (ranged from  $0.0014 \pm 0.001$   $\text{mgL}^{-1}$  to  $2.5714 \pm 0.980$   $\text{mgL}^{-1}$ ), nitrite+nitrate-nitrogen (ranged from  $0.0021 \pm 0.001$   $\text{mgL}^{-1}$  to  $0.8100 \pm 0.211$   $\text{mgL}^{-1}$ ) and total phosphorus (ranged from  $0.0712 \pm 0.024$   $\text{mgL}^{-1}$  to  $0.8670 \pm 0.3241$   $\text{mgL}^{-1}$ ) showed increasing trend from the coastal areas to the upstream. The highest chlorophyll *a* was recorded at Station 3 with mean values of  $10.27 \pm 0.84$   $\mu\text{gL}^{-1}$  and  $3.94 \pm 1.06$   $\mu\text{gL}^{-1}$  during high and low tide, respectively.

The mean zooplankton density was lowest in the coastal areas ( $16.21 \times 10^3 \pm 4679.07$  individuals  $\text{m}^{-3}$ ) and highest in Station 4 with a mean value  $119.81 \times 10^3 \pm 43338.73$  individuals  $\text{m}^{-3}$  during high tide. Copepods dominated zooplankton populations contributing >80% in all the stations throughout the sampling period. The other zooplankton groups such as cnidarians, appendicularians, polychaetes, ostracods, chaetognaths and shrimp larva were distributed in coastal estuarine areas whereas echinoderm larvae were restricted to coastal waters and cladocerans were found only at the upstream stations (Stations 3-5).

This study recorded 50 species of copepods which was dominated by the calanoids, *Acartia spinicauda*, *A. amboinensis* and *A. erythraea* accounting for 28%, 18% and 11% of the total copepod populations, respectively. High species diversity occurred in Station 1 decreasing towards the upstream. Species diversity varied depending on the tides, being higher during high tides and than low tides. Some copepod species were highly restricted to the high salinity levels (>30 PSU) while some species could tolerate wide range of salinity from 5 - >30 PSU. Major stenohaline species was *Oithona simplex* and major euryhaline species were *Acartia spinicauda*, *A. amboinensis* and *A. erythraea*. Zooplankton biomass study showed the highest in station 3 (dry weight  $612.30 \pm 26.31 \text{ mg m}^{-3}$  high tide) and lowest in Station 2 ( $150.05 \pm 54.71 \text{ mg m}^{-3}$  during low tide).

This study showed that water quality, nutrient concentrations and the distribution of zooplankton varied according to tides, where high tides resulted in better water quality, higher zooplankton density and higher species diversity compared to low tides. On the other hand, water quality was better and species diversity higher in coastal areas compared to other stations throughout the sampling period regardless of tides.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia  
sebagai memenuhi keperluan untuk ijazah Master Sains

**KESAN AIR PASANG SURUT KE ATAS PARAMETER FIZIKO-KIMIA,  
TABURAN TEMPORAL DAN SPATIAL ZOOPLANKTON DI  
KAWASAN KUALA SUNGAI LANGAT, MALAYSIA**

Oleh

**R.P. PRABATH KRISHANTHA JAYASINGHE**

**Jun 2005**

**Pengerusi: Profesor Fatimah Md. Yusoff, PhD**

**Fakulti: Sains**

Objektif kajian ini adalah untuk mengetahui kualiti air, taburan zooplankton serta struktur komuniti mengikut perubahan saliniti di sistem kuala Sungai Langat ( $2^{\circ} 46'N$ ,  $101^{\circ} 26' E$ ), semasa air pasang dan surut. Lima stesen persampelan telah dipilih bermula dari kawasan pesisiran hingga ke muara sungai. Stesen 1 terletak di kawasan pesisiran, stesen 5 terletak di bahagian hulu sungai sementara stesen lain (Stesen 2, 3 dan 4) terletak di antara kedua-dua kawasan tersebut merentasi gradien sungai. Zooplankton dan sampel air diambil dan dianalisis sekali dalam sebulan selama 12 bulan (Januari hingga Disember 2003).

Kualiti air, nutrien dan taburan zooplankton berubah berdasarkan stesen serta keadaan pasang surut. Min saliniti adalah antara  $32.94 \pm 1.20$  PSU di

kawasan pantai dan  $27.28 \pm 1.21$  PSU di stesen paling hulu semasa air pasang. Semasa air surut, min saliniti di stesen paling hulu sungai boleh turun sehingga  $1.00 \pm 0.90$  PSU. Nilai Kepekatan Oksigen Terlarut berada dalam lingkungan (min  $5.84 \pm 0.35$  hingga  $7.02 \pm 0.13$   $\text{mgL}^{-1}$ ) semasa air pasang dan menurun (min  $3.91 \pm 0.71$  hingga  $5.22 \pm 0.72$   $\text{mgL}^{-1}$ ) semasa air surut. Nutrien seperti jumlah ammonia- nitrogen (antara  $0.0014 \pm 0.001$   $\text{mgL}^{-1}$  hingga  $2.5714 \pm 0.980$   $\text{mgL}^{-1}$ ), nitrit+nitrat- nitrogen (antara  $0.0021 \pm 0.001$   $\text{mgL}^{-1}$  hingga  $0.8100 \pm 0.211$   $\text{mgL}^{-1}$ ) serta jumlah fosforus (antara  $0.0712 \pm 0.24$   $\text{mgL}^{-1}$  hingga  $0.8670 \pm 0.324$   $\text{mgL}^{-1}$ ) pula menunjukkan peningkatan kepekatan dari kawasan pesisiran hingga ke hulu sungai. Nilai klorofil *a* tertinggi yang direkodkan adalah di stesen 3 dengan nilai min masing masing  $10.27 \pm 0.84$   $\mu\text{gL}^{-1}$  dan  $3.94 \pm 1.06$   $\mu\text{gL}^{-1}$  semasa air pasang dan surut.

Min kepadatan zooplankton adalah paling rendah di kawasan persisiran pantai ( $16.21 \times 10^3 \pm 4679.07$  individu  $\text{m}^{-3}$ ) dan kepadatan adalah paling tinggi di kawasan menuju hulu sungai dengan nilai min  $119.81 \times 10^3 \pm 43338.73$  individu  $\text{m}^{-3}$  semasa air pasang. Populasi zooplankton Copepoda menyumbangkan >80% pada semua stesen sepanjang tempoh persampelan. Kumpulan zooplankton yang lain seperti cnidarian, appendikularian, polikete, ostrakoda, kaetognath dan larva udang didapati bertaburan di sepanjang persisiran muara sementara populasi ekinodermata bertumpu di sepanjang persisiran pantai dan kladosera hanya dijumpai pada stesen di hulu sungai (Stesen 3-5).



Kajian ini telah merekod sebanyak 50 spesies kopepod yang didominasi oleh kalanoid, *Acartia spinicauda*, *A. amboinensis* dan *A. erythrea* masing-masing sebanyak 28%, 18% dan 11% daripada jumlah populasi copepoda. Kepelbagaian spesies yang muncul di stesen 1 didapati menurun apabila menuju ke hulu sungai. Kepelbagaian spesies yang berbeza adalah bergantung kepada keadaan pasang surut di mana kepelbagaian adalah tinggi semasa air pasang dan rendah semasa air surut. Sebahagian spesies kopepoda tertumpu bertumpu kepada saliniti yang tinggi (>30 PSU) sementara sebahagian lagi spesies boleh bertoleransi dengan julat perbezaan saliniti yang luas iaitu daripada 5 PSU hingga 30 PSU bagi. Stenohalina species yang utama adalah *Oithona simplex* dan Eurihalina species yang utama pula terdiri daripada *Acartia spinicauda*, *A. amboinensis* dan *A. erythraea*. Kajian biojisim zooplankton adalah paling tinggi di stesen 3 (berat kering  $612.30 \pm 26.31 \text{ mg m}^{-3}$  semasa air pasang) dan paling rendah pada stesen 2 ( $150.05 \pm 54.71 \text{ mg m}^{-3}$  semasa air surut).

Kajian ini menunjukkan bahawa kualiti air, kepekatan nutrien dan taburan zooplankton adalah berbeza mengikut keadaan pasang surut, di mana pada keadaan air pasang, kualiti air didapati lebih baik, kepadatan zooplankton dan kepelbagaian zooplankton adalah lebih tinggi berbanding semasa air surut. Selain itu, kualiti air adalah baik dan kepelbagaian spesies adalah tinggi di kawasan persisiran berbanding dengan stesen lain sepanjang tempoh persampelan bergantung kepada sistem pasang surut.

## ACKNOWLEDGEMENTS

I am sincerely grateful to Her Excellency President of Democratic Socialist Republic of Sri Lanka, Chandrika Bandarnaike Kumaratunge for awarding me this scholarship from the President's Fund to pursue the Degree of Masters of Science in Universiti Putra Malaysia (UPM).

The Chairperson of my MS programme supervisory committee Professor Dr. Fatimah Md. Yusoff is greatly acknowledged for her invaluable advice, constructive criticisms, individual understanding and providing all the research facilities throughout the study. My deepest thanks and appreciation go to the members of the supervisory committee Associate Professor Dr. Aziz Arshad and Dr. Mahendran Shitan for their encouraging contribution.

My special thanks to Associate Professor Dr. U.S. Amarasinghe, University of Kelaniya Sri Lanka, for his invaluable advice and suggesting Professor Fatimah Md. Yusoff, UPM as a potential supervisor for my postgraduate studies. Professor Dr. Y. Ranga Reddy of Nagarjuna University, India, is highly appreciated for giving me the important technical knowledge on copepod taxonomy.

Financial support for the research work received from Malaysian Government Research Funds through Grant IRPA: 01-02-04-0392EA001, JICA



(Japan International Corporation Agency) and Japan Society for the Promotion of Science (JSPS). Head and other staff members of the Department of Biology, UPM are highly appreciated for providing necessary facilities and service to continue my study. Thanks to Marine Science Laboratory, Institute of Bioscience for providing microscopic facilities and Department of Mathematics for allowing me to use the computer packages.

I am grateful to Mr. Perumal Kuppan not only for giving all the support throughout the field and laboratory works but also sharing his experience and knowledge. I thank Mr. Kasim who was ready to bring us to the stormy waters in his fishing boat every month (throughout the sampling period). All my present and past laboratory mates Hamid, Said, Omid, Eduardo, Helen, Rozhan, Jamal, Nathrah, Ken, Adillah, Jimmy, Liza, Chin and Rita who gave their hand during sampling and/or data analysis. Without their pleasurable companionship, moral support and technical assistance, it would not be an easy task to complete this work within the time schedule. My heartfelt thanks go to Hazel for having a hard time to read the first draft of this thesis and her useful suggestions to improve the quality of the thesis.

My vocabulary is not wide enough to find the exact word to thank my friend Najim for giving me all the supports and helps whenever I needed specially the first few months of stay in Malaysia. I thank other Sri Lankan colleagues in Malaysia especially Dr. Chandana Duwe and Shantha for their company.

The officers in Sri Lankan High Commission for Malaysia helped me to receive scholarship money through Sri Lankan Government. My special appreciation goes to Nalanda Buddhist Society, Sri Serdang for their kind association with me during my stay in Malaysia.

Last but not least, my family members and close relatives are sincerely acknowledged for their understanding, love and everlasting moral support. Incomparable appreciation goes to my wife Mindi, for taking all the family responsibilities upon her shoulders. Finally, I express regret to my little daughter Nipulini, for my incapability to be with her during her early childhood.



I certify that an Examination Committee met on 29<sup>th</sup> June 2005 to conduct the final examination of R. P. Prabath Krishantha Jayasinge on his Master of Science thesis entitled “Tidal Influence on the Physico-Chemical Parameters, Temporal and Spatial Distribution of Zooplankton in Langat River-Estuarine Area, Malaysia” in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Jambari Ali, PhD**

Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Chairman)

**Japar Sidik Bujang, PhD**


Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Abdul Rahim Ismail, PhD**

Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Internal Examiner)

**Mazlan Abd. Ghaffar, PhD**

Associate Professor  
Faculty of Science and Technology  
Universiti Kebangsaan Malaysia  
(External Examiner)



**GULAM RUSUL RAHMAT ALI, PhD**  
Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia


Date: **22 AUG 2005**

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirements for the degree of Master of Science. The members of the Supervisory Committee are as follows:

**Fatimah Md. Yusoff, PhD**  
Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Aziz Arshad, PhD**  
Associate Professor  
Faculty of Science  
Universiti Putra Malaysia  
(Member)

**Mahendran Shitan, PhD**  
Lecturer  
Faculty of Science  
Universiti Putra Malaysia  
(Member)




---

**AINI IDERIS, PhD**  
Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date : 08 SEP 2005

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



**R.P. PRABATH KRISHANTHA JAYASINGHE**

**Date:** 02.06.2005

## TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENT	ix
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xvii
LIST OF FIGURES	xviii
 CHAPTERS	
I INTRODUCTION	1
Background of the study	1
Statement of the problem	2
Objectives	5
II LITERATURE REVIEW	6
Estuaries and water quality	6
Spatial and temporal distribution zooplankton	10
Composition of zooplankton	12
Biomass and abundance of zooplankton	15
Factors affecting zooplankton distribution	18
III GENERAL METHODOLOGY	21
Study area	21
Sampling sites	21
Field sampling	24
Physico- chemical parameters	24
Nutrient analysis	25
Total ammonia nitrogen (TAN)	25
Nitrate +nitrite nitrogen (NO <sub>3</sub> +NO <sub>2</sub> -N)	26
Total phosphorous (TP)	27
Chlorophyll <i>a</i>	28
Zooplankton studies	29
Sampling of zooplankton and preservation	29
Biomass estimates	30
Dry mass	30
Abundance of zooplankton	31
Data analysis	33
Physico-chemical parameters and nutrients	33
Biomass of zooplankton	33





	Density of zooplankton	33
IV	RESULTS	35
	Tide observation	35
	Variations of water quality parameters	36
	Zooplankton biomass	59
	Zooplankton abundance and composition	60
	Copepod composition	75
	Abundance of different copepod species	81
	Salinity ranges and occurrence of copepod species	91
	Species diversity (Shannon Index)	93
	Copepod group distribution (Cluster analysis)	94
V	DISCUSSION	97
	Tides and water quality	97
	Zooplankton community structure	101
	Copepod assemblages	107
VI	CONCLUSION AND RECOMMENDATIONS	114
	BIBLIOGRAPHY	116
	BIODATA OF THE AUTHOR	138

## List of Tables

Table	Page
1 The major estuaries in Asia and their length and basin area	7
2 Description of status of the tide at different sampling stations in Langat estuary during each sampling in through out the study period (January to December 2003)	35
3 Changes of different water quality parameters in different stations (random factor) and tide levels (fixed factor) over a 12-month study period based on two-way ANOVA analysis	38
4 Results of two-way ANOVA based on densities of total zooplankton and different major groups of zooplankton. The random factor is station (1-5) and the fixed factor is tide (high and low)	62
5 Pearson correlation coefficient (r) for zooplankton density and water quality parameters in Langat estuary	63
6 Percentage composition of zooplankton (mean $\pm$ SE) at different stations during high and low tides in Langat estuary	64
7 List of species of copepods and mean salinity ranges recorded in Langat estuary during 2003	77
8 Mean percentage distribution of most abundant copepod species (>1% any station) at different stations during high and low tides in Langat estuary during 2003	80
9 Number of copepod families, genera and species recorded in Langat estuary during 2003	81
10 Changes of different copepod species at different stations (random factor) and tide levels (fixed factor) over a 12 month period based on two-way ANOVA analysis	84



## List of Figures

Figure		Page
1	Langat estuary showing sampling sites (Stations 1-5)	22
2	Variations of water temperature in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	37
3	Means of water temperature in Langat estuary during high tide (A) and low tide (B)	39
4	Variations of salinity in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	40
5	Means salinity in Langat estuary during high tide (A) and low tide (B)	41
6	Means of Secchi disk depth in Langat estuary during high tide (A) and low tide (B)	43
7	Variations of dissolved oxygen (DO) in Langat estuary at different sampling stations during the months of high tide (A) and low tide (B) in 2003	45
8	Means of dissolved oxygen (DO) in Langat estuary during high tide (A) and low tide (B)	46
9	Variations of pH value in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	47
10	Means of pH values in Langat estuary during high tide (A) and low tide (B)	48
11	Variations of total ammonia nitrogen (TAN) in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	50
12	Means of total ammonia nitrogen (TAN) in Langat estuary during high tide (A) and low tide (B)	51
13	Variations of nitrate+nitrite-nitrogen ( $\text{NO}_3+\text{NO}_2\text{-N}$ ) in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	52
14	Means of nitrate+nitrite nitrogen ( $\text{NO}_3+\text{NO}_2\text{-N}$ ) in Langat estuary during high tide (A) and low tide (B)	53
15	Variations of total phosphorus (TP) in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	55
16	Means of total phosphorus (TP) in Langat estuary during high tide (A) and low tide (B)	56
17	Variations of chlorophyll <i>a</i> in Langat estuary at different stations during the months of high tide (A) and low tide (B) in 2003	57

18	Means of chlorophyll <i>a</i> in Langat estuary during high tide (A) and low tide (B)	58
19	Mean valules of zooplankton biomass (dry weight mgm <sup>-3</sup> ) in Langat estuary during high tide (H) and low tide (L) at different sampling stations during 2003	59
20	Mean density ( $\pm$ SE) of total zooplankton ( $\times 10^3$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	61
21	Mean density ( $\pm$ SE) of copepods ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide (A) and low tide (B)	65
22	Mean density ( $\pm$ SE) of nauplii ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	67
23	Mean density ( $\pm$ SE) of cnidarians ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	68
24	Mean density ( $\pm$ SE) of appendicularians ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide (A) and low tide (B)	70
25	Mean density ( $\pm$ SE) of cheatognaths ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide (A) and low tide (B)	71
26	Mean density ( $\pm$ SE) of polychaetes ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide (A) and low tide (B)	72
27	Mean density ( $\pm$ SE) of echinoderm larvae ( $\times 10^2$ individuals m <sup>-3</sup> ) during high tide (A) and cladocerans ( $\times 10^2$ individuals m <sup>-3</sup> ) during low tide (B) in Langat estuary	74
28	Mean density ( $\pm$ SE) of <i>Acartia spinicauda</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	83
29	Mean density ( $\pm$ SE) of <i>Acartia pacifica</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	85
30	Mean density ( $\pm$ SE) of <i>Acartia erythraea</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	86
31	Mean density ( $\pm$ SE) of <i>Paracalanus aculeatus</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	88
32	Mean density ( $\pm$ SE) of <i>paracalanus crassirostris</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	89
33	Mean density ( $\pm$ SE) of <i>Parvocalanus denudatus</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	90
34	Mean density ( $\pm$ SE) of <i>Bestiolina similix</i> ( $\times 10^2$ individuals m <sup>-3</sup> ) in Langat estuary during high tide and low tide (B)	92
35	Mean values of species diversity index (Shannon Index) at different stations during high tide (HT) and low tide (LT) in Langat estuary during 2003	93
36	Dendrogram of copepod group classification during high tide (most abundant copepod species were used for the analysis) in Langat estuary during 2003	94

37	Dendrogram of copepod group classification during low tide (most abundant copepod species were used for the analysis) in Langat estuary during 2003	96
----	---	----

## CHAPTER I

### INTRODUCTION

#### Background of the study

Zooplankton is a very important group of animals in aquatic ecosystems because they are the basis for food web. They constitute the important intermediate steps in the food pyramid. They transfer the organic energy produced by unicellular algae through photosynthesis to higher trophic levels such as pelagic fish stocks. Their reproductive cycles, growth, reproduction and survival rates are all important factors influencing recruitment of fish stocks (Harris *et al.*, 2000). Furthermore, zooplankton play critical role in the food intake of many invertebrates and reef fishes (Emery, 1968; Hammer and Carleton, 1979; Alldredge and King, 1980; Robichaux *et al.*, 1981; Noda *et al.*, 1998) and corals (Hammer *et al.*, 1988). Many small sized zooplankton are the main food source of carnivorous zooplanktors (Moore and Sander, 1979).

In Malaysia, marine zooplankton studies were carried out in 1928 (Keller and Richads, 1967), followed by Sewell (1933); Tham (1953); Pathansali (1968) and Tham *et al.* (1970). After that, the main focus of marine zooplankton research has gone to the Straits of Malacca (Chua and Chong, 1975; Arvin, 1977; Idris *et al.*, 2000; Johan *et al.*, 2000; Rezai, 2002; Rezai *et al.*, 2003a, 2003b, 2004a,

2004b) due to its economical, ecological and biological importance (Rezai 2002). Only a few zooplankton studies were carried out in the South China Sea (Chark and Saufi, 1987; Othman, 1988), East Coast of Malaysia (Chua, 1980; Jivaluk, 1999a) and Sabah and Sarawak coastal waters of Malaysia (Jivaluk, 1999b). Several authors have described different aspects of freshwater zooplankton in Malaysia such as copepods (Lai and Fernando, 1978) and cladocerans (Idris, 1983). These studies were mainly on species composition (Fernando, 1980a; 1980b), species distribution (Fernando, 1980c) and ecology (Lim *et al.*, 1984). Eventhough Malaysia has a vast number of estuaries, no zooplankton studies were done in those waters. Estuaries are one of the most productive ecosystems (Odum, 1971). They are important both ecologically and economically as they serve as breeding, feeding and nursery grounds for many aquatic organisms including fish (Ross and Epperly, 1985; Deegan and Day, 1985, 1986) and shrimp species (Meager *et al.*, 2003). Furthermore, estuarine ecosystems support the wildlife (Hock and Sasekumar, 1979; Smith and Odum, 1981) such as migratory birds and wild fowls.

## Statement of the Problem

Estuaries represent areas of prime interest for human activities such as navigation (Carlberg, 1980), domestic and industrial garbage disposal (Carlberg, 1980; Chau, 1999; Kress *et al.*, 2002), human settlement (Day *et al.*, 1989; Wang *et al.*, 2004), fisheries and aquaculture (Jennerjahn *et al.*, 2004) and recreational activities (Baird *et al.*, 1986; Costanza *et al.*, 1997). These human activities together like deforestation, intensive agriculture, livestock farming, sand mining, river diversion and conversion of the estuarine mangroves to aquaculture ponds may alter the original structure of the estuaries and marine environment (Morton and Blackmore, 2001; Jennerjahn *et al.*, 2004). Therefore, most estuaries are threatened by human activities.

Like other estuaries in the world, estuarine areas in Malaysia have been subjected to strong anthropogenic impacts due to massive aquaculture, agriculture and siltation due to soil erosion from land base activities (Chan, 1985; Abdullah, 1995). Almost 80% of the estuaries and rivers in Malaysia are polluted (Law and Mohammad Moshin, 1980; Chye and Furtado, 1982). Many anthropogenic activities such as massive use of nutrient-rich products, especially agro-industries, have dramatically enhanced eutrophication by increasing the nutrient input into water-bodies, especially estuaries (Abdullah, 1995). Although estuaries and coastal areas in Malaysia are fast changing due to pollution, scientific studies in these areas are lacking. Deleterious effects caused by pollutants may not be remarkable in the initial



stages as they are masked by dilution, especially during the high tide. However, if no effective steps are taken to reduce the pollution, eventually changes in water quality will result not only in reduced biodiversity and other undesirable ecological impacts but also creates social and economical problems. Furthermore, this will also generate problems in human health (Cato *et al.*, 1980). At the end, great costs are involved if decisions are made to clean up polluted areas and these manipulations will follow by the establishing the original ecosystem. (Cato *et al.*, 1980). Therefore, identification and quantification of the level of pollution should form an important part of managing both land and water resources within a particular river catchment (Petts and Calow, 1996).

As one of the first steps towards managing the estuarine environment, basic data on physical-chemical characteristics and biological entities are required. Similar to other ecological studies, information on zooplankton populations along river-estuaries in Malaysia is deficient. The present study focused on the temporal and spatial distribution of zooplankton in Langat river-estuarine, Malaysia, giving emphasis to its relationship with water quality parameters and nutrients. This study is important, as part of conservation of the biodiversity and results will be helpful to take sustainable management issues to reduce the pollution effects.